HEDIS® & QARR Provider Reference Guide - Measurement Year 2023





Table of Contents

Glossary of Terms:	
Effectiveness of Care- Prevention and Screening Measures:	
Weight Assessment & Counseling for Nutrition and Physical Activity for Children & Adolescents (WCC)	
Childhood Immunizations Status (CIS)	
Immunizations for Adolescents (IMA)	
Lead Screening in Children (LSC)	
Breast Cancer Screening (BCS)	
Cervical Cancer Screening (CCS)	8
Colorectal Cancer Screening (COL)	
Chlamydia Screening in Women (CHL)	
Care of Older Adults (COA)	9
Oral Evaluation, Dental Services (OED)	9
Topical Fluoride for Children (TFC)	
Medical Assistance with Smoking and Tobacco Cessation (MSC)	9
COVID-19 Immunization Status (CVS)	9
Effectiveness of Care- Respiratory Conditions Measures:	10
Appropriate Testing for Pharyngitis (CWP)	10
Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR)	10
Pharmacotherapy Management of COPD Exacerbation (SPC)	10
Asthma Medication Ratio (AMR)	1
Effectiveness of Care- Cardiovascular Conditions Measures:	12
Controlling High Blood Pressure (CBP)	12

Persistence of Beta-Blocker Treatment After a Heart Attack (PBH)	12
Statin Therapy for Patients with Cardiovascular Disease (SPC)	13
Cardiac Rehabilitation (CRE)	14
Effectiveness of Care- Diabetes Measures:	15
Hemoglobin A1c Control for Patients with Diabetes (HBD)	15
Blood Pressure Control for Patients with Diabetes (BPD)	15
Eye Exam for Patients with Diabetes (EED)	15
Kidney Health Evaluation for Patients with Diabetes (KED)	16
Statin Therapy for Patients with Diabetes (SPD)	16
Effectiveness of Care- Musculoskeletal Conditions Measures:	18
Osteoporosis Management in Women Who Had a Fracture (OMW)	18
Osteoporosis Screening in Older Women (OSW)	18
Effectiveness of Care- Behavioral Health Measures:	19
Diagnoses of Mental Health Disorders (DMH)	19
Antidepressant Medication Management (AMM)	19
Follow-up Care for Children Prescribed ADHD Medication (ADD)	19
Follow-Up After Hospitalization for Mental Illness (FUH)	19
Follow-Up After Emergency Department Visit for Mental Illness (FUM)	20
Diagnosed Substance Use Disorders (DSU)	20
Follow-Up After High-Intensity Care for Substance Use Disorder (FUI)	20
Follow-Up After Emergency Department Visit for Substance Use (FUA)	20
Pharmacotherapy for Opioid Use Disorder (POD)	21
Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD)	21
Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD)	21

Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC)(SMC)	22
Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA)	22
Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)	22
Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults (DMS-E)	22
Depression Remission or Response for Adolescents and Adults (DRR-E)	23
Effectiveness of Care- Care Coordination Measures:	24
Advance Care Planning (ACP)	24
Transitions of Care (TRC)	24
Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic Conditions (FMC)	24
HIV Care	25
Viral Load Suppression	25
Sexually Transmitted Diseases: Chlamydia Screening	25
Sexually Transmitted Diseases: Gonorrhea Screening	25
Sexually Transmitted Diseases: Syphilis Screening	25
Health Plan Descriptive Information	26
Social Need Screening and Intervention (SNS-E)	26
Language Diversity of Membership (LDM)	26
Race/Ethnicity Diversity of Membership (RDM)	26
Access/Availability of Care Measures:	27
Adults' Access to Preventative/Ambulatory Health Services (AAP)	27
Initiation and Engagement of Substance Use Disorder Treatment (IET)	27
Prenatal and Postpartum Care (PPC)	27
Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP)	27
Measures Collected Through the Health Outcomes Survey:	28

Medicare Health Outcomes Survey (HOS)	28
Fall Risk Management (FRM)	28
Management of Urinary Incontinence in Older Adults (MUI)	28
Physical Activity in Older Adults (PAO)	28
Utilization and Risk Adjusted Utilization Measures:	
Well-Child Visits in the First 30 Months of Life (W30)	29
Child and Adolescent Well-Care Visits (WCV)	29
Plan All-Cause Readmissions (PCR)	29
Hospitalization Following Discharge from a Skilled Nursing Facility (HFS)	29
Acute Hospitalization Utilization (AHU)	
Emergency Department Utilization (EDU)	30
Hospitalization for Potentially Preventable Complications (HPC)	30
Emergency Department Visits for Hypoglycemia in Older Adults with Diabetes (EDH)	30
References and Comments:	31

Healthcare Effectiveness Data and Information Set (HEDIS®) & Quality Assurance Reporting Requirements (QARR) Provider Reference Guide - MY2023

Glossary of Terms:

Measure Name: The name of the HEDIS® measure used for reporting

Measure Code: The code used to identify the measure name

Age Criteria: The age of the population that is eligible for the measure

Measure Description: The description of what is being measured in each domain

ICD10 Codes: The diagnosis code assigned to conditions and diseases used for claims reporting

CPT Codes: Codes used to describe the treatment being provided

Medication List: List of medications from the current year HEDIS® specifications that meet numerator compliance for the attributed measure

HCPCS Codes: Codes used to describe supplies, equipment and devices being provided

LOINC Codes: Codes use to describe the tests being provided, such as laboratory tests and vital signs

Exclusions: Some measures remove eligible members from a measure they are otherwise eligible for (ex: hospice, palliative care, ESRD/dialysis)

Measurement Year (MY): Refers to the current calendar year (January 1, 2023 – December 31, 2023), for the measures unless otherwise specified

Effectiveness of Care- *Prevention and Screening Measures:*

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Weight Assessment & Counseling for Nutrition and Physical Activity for Children & Adolescents (WCC)	3 to 17 years of age	Eligible members with an outpatient PCP and/or OB/GYN visit with documentation of BMI profile and counseling for nutrition and physical activity, within the MY.	ICD 10 Codes: Z68.51-Z68.54, Z71.3 CPT Codes: 97802-97804 Exclusion: Members on/using hospice services (99377); members with a pregnancy diagnosis during the MY, and members who expired during the MY.	SelectHealth
Childhood Immunizations Status (CIS)	2 years of age	Children who have received four doses of DTaP (diphtheria, tetanus, and pertussis) and PCV (pneumococcal conjugate); three doses of IPV (polio), HepB (hepatitis B), and HIB (haemophilus influenza B), two to three doses of RV (rotavirus), two Flu vaccines (influenza), and one dose of MMR (measles, mumps, and rubella), VZV (chicken pox) and HepA (Hepatitis A) by their second birthday.	ICD 10 Codes: NA CPT Codes: 90697, 90670, 90644, 90680/90681, 90655, 90674, 90707, 90710, 90633 Exclusion: Members on/using hospice services (99377), members who expired during the MY, diagnosis of HIV (B20, B97.35) and immunocompromised conditions (D84.9).	SelectHealth
Immunizations for Adolescents (IMA)	13 years of age	Adolescents who have received one dose of Tdap (tetanus, diphtheria, and pertussis) and meningococcal vaccines, and have had the HPV (human papillomavirus) vaccine series finalized, by their 13 th birthday.	ICD 10 Codes: NA CPT Codes: 90649, 90619, 90715 Exclusions: Members on/using hospice services (99377) and members who expired during the MY.	SelectHealth
Lead Screening in Children (LSC)	2 years of age	Children who have completed at least one lead blood test (capillary or venous) by their second birthday, to determine lead poisoning.	ICD 10 Codes: NA CPT Codes: 83655 LOINC Codes: 5671-3 Exclusions: Members on/using hospice services (99377) and members who expired during the MY.	SelectHealth
Breast Cancer Screening (BCS)	Women 52 to 74 years of age	Women between 52 to 74 years of age, who have had a mammogram October 1 st two	ICD 10 Codes: NA CPT Codes: 77061-77063, 77065-77067	SelectHealth, Total, EasyCare

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
		years prior through the end of the MY, to screen for breast cancer. *NCQA implemented an electronic version of this measure (BCS-E). Providers are encouraged to submit data administratively.	Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who have expired during the MY, and those who have either a unilateral (19180) or bilateral mastectomy (Z90.13) modifier (optional), members 66 and older residing in a facility or members 66 and older with a diagnosis frailty (L89.000) <i>and</i> advanced illness (A81.00).	and EasyCare Plus
Cervical Cancer Screening (CCS)	Women 21 to 64 years of age	Women 21 to 64 years of age, who have had a cervical cytology (21 to 64) within 3 years, cervical high-risk human papillomavirus (30-64) within 5 years or both cytology/hrHPV cotesting (30-64) within the last 5 years, to screen for cervical cancer. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data administratively.	ICD 10 Codes: NA CPT Codes: 88141-88143, 87624, 87625 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, or have had a hysterectomy (57530) and/or absence of cervix (Q51.5) (optional).	SelectHealth
Colorectal Cancer Screening (COL)	45 to 75 years of age	Members 45 to 75 years of age, who have had at least one of the following cancer screenings within the MY and prior year: Fecal Occult Blood Test (FOBT), Flexible Sigmoidoscopy, Colonoscopy, CT Colonography and/or Stool DNA with FIT testing.	ICD 10 Codes: NA CPT Codes: 82270, 45330, 44388,74261, 81528 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 and older residing in a facility or members 66 and older with a diagnosis frailty (L89.000) and advanced illness (A81.00), and members with a diagnosis of colorectal cancer (C18.0) or total colectomy (44150) (optional).	SelectHealth, Total, EasyCare and EasyCare Plus
Chlamydia Screening in Women (CHL)	Women 16 to 24 years of age	Sexually active women between 16 to 24 years of age, who have at least one test for chlamydia within the MY.	ICD 10 Codes: NA CPT Codes: 87110, 87320, 87490-87492	SelectHealth

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
			Exclusions: Members on/using hospice services (99377), and members who have expired during the MY.	
Care of Older Adults (COA)	66 years of age	Members who are 66 years of age who have completed all three at least once within the MY- Pain and functional status assessments, as well as medication review (including OTC and supplements).	ICD 10 Codes: NA CPT Codes: 99483, 90863, 99495 HCPCS Codes: G8427, G0438 Exclusions: Members on/using hospice services (99377), members who expired during the MY, and services provided in an acute inpatient setting (99221).	Total, and EasyCare Plus
Oral Evaluation, Dental Services (OED)	Members under 21 years of age	Members under 21 years of age, who have completed a comprehensive or periodic oral exam with a dental provider during the MY.	ICD 10 Codes: NA CPT Codes: NA Exclusions: Members on/using hospice services (99377), or members who have expired during the MY.	SelectHealth
Topical Fluoride for Children (TFC)	1 to 4 years of age	Members between 1 and 4 years of age, who have had two or more fluoride varnish applications completed during the MY.	ICD 10 Codes: NA CPT Codes: 99188 Exclusions: Members on/using hospice services (99377), or members who have expired during the MY.	SelectHealth
Medical Assistance with Smoking and Tobacco Cessation (MSC) *CAHPS	18 years of age and older	This is a patient survey based measure.	ICD 10 Codes: NA CPT Codes: NA Exclusions: NA	SelectHealth, Total, EasyCare and EasyCare Plus
COVID-19 Immunization Status (CVS)	6 months of age and older	Members 6 months of age and older, who have received the primary vaccine series, and members 5 years of age and older, who have received both the primary vaccine series and a booster.	ICD 10 Codes: NA CPT Codes: 91308, 91307, 91303 Exclusions: Dually enrolled members in Medicaid and Medicare during the MY.	SelectHealth

Effectiveness of Care- *Respiratory Conditions Measures:*

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Appropriate Testing for Pharyngitis	3 years of age and	Members 3 years of age and older, who were evaluated and diagnosed with pharyngitis,	ICD 10 Codes: J02.0, J02.8, J03.01 CPT Codes: 87070, 87081, 87430	SelectHealth, Total,
(CWP)	older	given an antibiotic and received a strep test	Exclusions: Members on/using hospice services	EasyCare
,		(group A streptococcus).	(99377), and members who expired during the	and
			MY.	EasyCare Plus
Use of Spirometry	42 years of	Members 40 years of age or older, who	ICD 10 Codes: J44.0, J44.1, J.44.9	SelectHealth,
Testing in the	age or older	received adequate spirometry testing to	CPT Codes: 94010, 94014-94016, 94060, 94375	Total,
Assessment and		determine a new diagnosis of COPD, between	Exclusions: Members on/using hospice services	EasyCare
Diagnosis of COPD		July 1 of prior year through June 30 of the	(99377), and members who expired during the	and
(SPR)		MY.	MY.	EasyCare
				Plus
Pharmacotherapy	40 years of	Members 40 years of age, which were	ICD 10 Codes: J44.0, J44.9, J43.0, J43.9	SelectHealth,
Management of	age	hospitalized or had an emergency room visit	Medication List: Cortisone, Dexamethasone,	Total,
COPD Exacerbation		between January 1 st and November 30 th , with	Hydrocortisone, Methylprednisolone,	EasyCare
(PCE)		COPD exacerbation, who were prescribed	Prednisolone, Prednisone, Aclidinium bromide,	and
		appropriate medications (a systemic	Ipratropium, Tiotropium, Umeclidinium,	EasyCare
		corticosteroid or a bronchodilator).	Albuterol, Aformoterol, Formoterol,	Plus
			Indacaterol, Levalbuterol, Metaproterenol, Olodaterol, Salmeterol, Albuterol-ipratropium,	
			Budesonide-formoterol, Fluticasone-salmeterol,	
			Fluticasone-vilanterol, Fluticasone furoate-	
			umeclidinium-vilanterol, Formoterol-aclidinium,	
			Formoterol-glycopyrrolate, Formoterol-	
			mometasone, Glycopyrrolate-indacaterol,	
			Olodaterol-tiotropium and Umeclidinium-	
			vilanterol.	
			Exclusions: Members on/using hospice services	
			(99377), members who have expired during the	
			MY, and nonacute hospitalizations.	

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Asthma Medication Ratio (AMR)	5 to 64 years of age	Members 5 to 64 years of age, with a diagnosis of persistent asthma that have a controller medications vs total asthma medications prescribed ratio of at least 0.50.	ICD 10 Codes: J45.21, J45.51, J45.991 Medication List: Omalizumab, Dupilumab, Benralizumab, Mepolizumab, Reslizumab, Budesonide-formoterol, Fluticasone-salmeterol, Fluticasone-vilanterol, Formoterol- mometasone, Beclomethasone, Budesonide, Ciclesonide, Flunisolide, Fluticasone, Mometasone, Montelukast, Zafirlukast, Zileuton and Theophylline. Exclusions: Members on/using hospice services (99377), members who have expired during the MY, members who had a diagnosis of emphysema (J43.0), COPD (J44.0), obstructive chronic bronchitis, chronic respiratory conditions (J68.4), cystic fibrosis and acute respiratory failure (J96.00), and members without any asthma controlling medications dispensed during the MY.	SelectHealth

Effectiveness of Care- *Cardiovascular Conditions Measures:*

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Controlling High	18 to 85	Members 18 to 85 years of age with a	ICD 10 Codes: I10	SelectHealth,
Blood Pressure	years of age	diagnosis of HTN (hypertension) whose most	CPTII Codes: 3079F, 3078F, 3080F, 3074F,	Total,
(CBP)		recent blood pressure measurements, during	3075F, 3077F	EasyCare
		the MY, show an adequate control of	Exclusions: Members on/using hospice services	and
		condition, as evidenced by results of	(99377) or palliative care (Z51.5), members who	EasyCare
		<140/90mm Hg.	have expired during the MY, members 66 years	Plus
			of age or older residing in a facility, members 66	
			to 80 years of age with a frailty (L89.000) and	
			advanced illness (A81.00) diagnosis, and	
			members 81 and older with at least two	
			indications of frailty (L89.000) during the MY,	
			members with a diagnosis of End-stage renal	
			disease (ESRD) (N18.5), nephrectomy (50220) or	
			kidney transplant (50360), and female members	
			who are pregnant (O00.0).	
Persistence of Beta-	18 years of	Members 18 years of age and older who were	ICD 10 Codes: I21.01, I21.19, I21.29	SelectHealth,
Blocker Treatment	age and	discharged from a hospitalization for AMI	Medication List: Carvedilol, Labetalol, Nadolol,	Total,
After a Heart Attack	older	(acute myocardial infarction), that were	Pindolol, Propranolol, Timolol, Sotalol,	EasyCare
(PBH)		prescribed a six-month treatment of	Acebutolol, Atenolol, Betaxolol, Bisoprolol,	and
		persistent beta-blockers.	Metorpolol, Nebivolol, Atenolol-chlorthalidone,	EasyCare
			Bendroflumethiazide-nadolol, Bisoprolol-	Plus
			hydrochlorothiazide, Hydrochlorothiazide-	
			metroprolol and Hydrochlorothiazide-	
			propranolol.	
			Exclusions: Members on/using hospice services	
			(99377), members who expired during the MY,	
			members 66 years of age or older residing in a	
			facility, members 66 to 80 years of age with a	
			frailty (L89.000) <i>and</i> advanced illness (A81.00)	
			diagnosis, and members 81 and older with a	

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Statin Therapy for Patients with Cardiovascular Disease (SPC)	Males 21 to 75 years of age; Females 40 to 75 years of age	Male members between 21 to 75 years of age, and female members between 40 to 75 years of age, who have had an ASCVD diagnosis (atherosclerotic cardiovascular disease) and has received statin therapy at least once during the MY, as well as maintained an 80% or higher of statin adherence.	frailty (L89.000) diagnosis, members diagnosed with asthma (J45.21), COPD (J44.0), obstructive chronic bronchitis, chronic respiratory conditions (J68.4) and hypotension, members who have intolerance or allergic to betablockers, and members who have been dispensed an indicative asthma medication. ICD 10 Codes: I21.01, I21.9, I25.2 High-intensity Statin Medication List: Atorvastatin 40-80mg, Amlodipine-atorvastatin 40-80mg, Rosuvastatin 20-40mg, Simvastatin 80mg and Ezetimibe-simvastatin 80mg. Moderate-intensity Statin Medication List: Atorvastatin 10-20mg, Amlodipine-atorvastatin 10-20mg, Rosuvastatin 5-10mg, Simvastatin 20-40mg, Ezetimibe-simvastatin 20-40mg, Pravastatin 40-80mg, Lovastatin 40mg, Fluvastatin 40-80mg and Pitavastatin 1-4mg. Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 years of age or older residing in a facility or have a diagnosis of frailty (L89.000) and advanced illness (A81.00), members diagnosed with ESRD (N18.5), cirrhosis (K70.30), myalgia (M79.10), myositis (M60.80), myopathy (G72.9) or rhabdomyolysis (M62.82) and members with a pregnancy (O00.0) diagnosis, or received IVF (in	SelectHealth, Total, EasyCare and EasyCare Plus
			vitro fertilization) (HCPCS S4015) or prescribed clomiphene during the MY or the year prior.	

Measure Name	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of
(Measure Code)				Business
				(LOB)
Cardiac	18 years of	Members 18 years of age and older, who	ICD 10 Codes: I21.01, I21.9, I25.2	SelectHealth,
Rehabilitation (CRE)	age and	participated in cardiac rehabilitation	CPT Codes: 93797, 93798	Total,
	older	following an eligible cardiac event (such as	Exclusions: Members on/using hospice services	EasyCare
		MI, CABG or heart transplant).	(99377) or palliative care (Z51.5), members who	and
		Rates reported include initiation (2 or more	expired during the MY, members 66 years of	EasyCare
		sessions within 30 days), engagement 1 (12 or	age or older residing in a facility, members 66 to	Plus
		more sessions within 90 days), engagement 2	80 years of age with a frailty (L89.000) and	
		(24 or more sessions within 180 days) and	advanced illness (A81.00) diagnosis, members	
		achievement (36 or more sessions within 180	81 and older with two or more indications of	
		days).	frailty (L89.000). Members diagnosed with an	
			MI (I21.01), CABG (33510), heart or heart &	
			lung transplant (33927) or heart valve	
			repair/replacement (33361) within 180 days	
			after initial event are also excluded.	

Effectiveness of Care- *Diabetes Measures:*

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Hemoglobin A1c	18 to 75	Members between 18 to 75 years of age,	ICD 10 Codes: E10.10, E10.36, E11.00	SelectHealth,
Control for Patients	years of age	diagnosed with diabetes (types 1 and 2), who	CPT Codes: 83036, 83067	Total,
with Diabetes (HBD)		have a completed hemoglobin A1c within the	CPTII Codes: 3044F, 3051F, 3052F, 3046F	EasyCare
		MY.	Exclusions: Members on/using hospice services	and
		A1c levels <8.0 indicate good control and	(99377) or palliative care (Z51.5), members who	EasyCare
		levels >9.0 indicate poor control.	have expired during the MY, members 66 years	Plus
			of age or older residing in a facility, members 66	
			or older with a frailty (L89.000) and advanced	
			illness (A81.00) diagnosis, members without a	
			diagnosis of diabetes and diagnosed with	
			polycystic ovarian syndrome (E28.2), gestational	
			(O24.419) or steroid induced diabetes (E09.9).	
Blood Pressure	18 to 75	Members between 18 to 75 years of age,	ICD 10 Codes: E10.10, E10.36, E11.00	SelectHealth,
Control for Patients	years of age	diagnosed with diabetes (types 1 and 2),	CPTII Codes: 3079F, 3078F, 3080F, 3074F,	Total,
with Diabetes (BPD)		whose most recent blood pressure	3075F, 3077F	EasyCare
		measurements, during the MY, show an	Exclusions: Members on/using hospice services	and
		adequate control of condition, as evidenced	(99377) or palliative care (Z51.5), members who	EasyCare
		by results of <140/90mm Hg.	expired during the MY, members 66 years of	Plus
			age or older residing in a facility, members 66	
			and older with a frailty (L89.000) and advanced	
			illness (A81.00) diagnosis, members without a	
			diagnosis of diabetes and diagnosed with	
			polycystic ovarian syndrome (E28.2), gestational	
			(O24.419) or steroid (E09.9) induced diabetes.	
Eye Exam for	18 to 75	Members between 18 to 75 years of age,	ICD 10 Codes: E10.10, E10.36, E11.00	SelectHealth,
Patients with	years of age	diagnosed with diabetes (types 1 and 2), who	CPT Codes: 67028, 67030, 92201, 92230	Total,
Diabetes (EED)		have completed a retinal eye exam.	CPTII Codes: 2022F (evidence of retinopathy),	EasyCare
			2023F (no evidence of retinopathy), 3072F (no	and
			evidence of retinopathy in the prior year)	EasyCare
				Plus

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Kidney Health Evaluation for Patients with Diabetes (KED)	18 to 85 years of age	Members between 18 to 85 years of age, diagnosed with diabetes (types 1 and 2), who completed the eGFR (estimated glomerular filtration rate) and uACR (urine albumincreatinine ratio) lab tests during the MY. *The uACR must have both a quantitative urine albumin test and a urine creatinine test to be complete.	Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 years of age or older residing in a facility, members 66 and older with a frailty (L89.000) and advanced illness (A81.00) diagnosis, members without a diagnosis of diabetes and diagnosed with polycystic ovarian syndrome (E28.2), gestational (O24.419) or steroid (E09.9) induced diabetes. ICD 10 Codes: E10.10, E10.36, E11.00 CPT Codes: 82043, 82570, 80047 LOINC Codes: 13705-9, 48642-3, 1754-1, 2161-8 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) and advanced illness (A81.00) diagnosis, members 81 and older with two indications of frailty (L89.000) diagnosis, members diagnosed with ESRD (N18.5) or on dialysis, and members without a diagnosis of diabetes and diagnosed with polycystic ovarian syndrome (E28.2), gestational (O24.419) or steroid (E09.9) induced diabetes.	SelectHealth, Total, EasyCare and EasyCare Plus
Statin Therapy for Patients with Diabetes (SPD)	40 to 75 years of age	Members between 40 to 75 years of age, with a diagnosis of diabetes, who do not have an ASCVD diagnosis (atherosclerotic cardiovascular disease), and has received statin therapy, as well as maintained an 80% of statin adherence.	ICD 10 Codes: E10.10, E10.36, E11.00 High-intensity Statin Medication List: Atorvastatin 40-80mg, Amlodipine-atorvastatin 40-80mg, Rosuvastatin 20-40mg, Simvastatin 80mg and Ezetimibe-simvastatin 80mg.	SelectHealth, Total, EasyCare and EasyCare Plus

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
			Moderate-intensity Statin Medication List:	
			Atorvastatin 10-20mg, Amlodipine-atorvastatin	
			10-20mg, Rosuvastatin 5-10mg, Simvastatin 20-	
			40mg, Ezetimibe-simvastatin 20-40mg,	
			Pravastatin 20-40mg, Lovastatin 40mg,	
			Fluvastatin 40-80mg and Pitavastatin 1-4mg.	
			Low-intensity Statin Medication List:	
			Ezetimibe-simvastatin 10mg, Fluvastatin 20mg,	
			Lovastatin 10-20mg, Pravastatin 10-20mg and	
			Simvastatin 5-10mg.	
			Exclusions: Members on/using hospice services	
			(99377) or palliative care (Z51.5), members who	
			expired during the MY, members 66 years of	
			age or older residing in a facility, members 66	
			and older with a frailty (L89.000) and advanced	
			illness (A81.00) diagnosis, members diagnosed	
			with MI (I21.9), CABG or PCI during the MY, and	
			members with a diagnosis of ESRD (N18.5),	
			cirrhosis (K70.30), myalgia (M79.1), myositis	
			(M60.80), myopathy (G72.9) or rhabdomyolysis	
			(M62.82) and members with a pregnancy	
			(O00.0) diagnosis, or received IVF (in vitro	
			fertilization) (HCPCS S4015) or prescribed	
			clomiphene during the MY and members	
			without a diagnosis of diabetes and diagnosed	
			with polycystic ovarian syndrome (E28.2),	
			gestational (O24.419) or steroid (E09.9) induced	
			diabetes.	

Effectiveness of Care- *Musculoskeletal Conditions Measures:*

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Osteoporosis Management in Women Who Had a Fracture (OMW)	67 to 85 years of age	Female members between 67 to 85 years of age, who have had a BMD (bone mineral density) test or prescribed a medication for osteoporosis treatment within six months, after suffering a fracture.	ICD 10 Codes: M80.00XA, M80.80XA CPT Codes: 76977, 77078, 77080-77081 HCPCS Codes: J0897, J1740, J3110, J3111, J3489 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 67 years of age or older residing in a facility, members 67 to 80 years of age with a frailty (L89.000) and advanced illness (A81.00) diagnosis, members 81 and older with two or more indications of frailty (L89.000) diagnosis, and members with a BMD test within 2 years prior to fracture event, had encounters for osteoporosis therapy or claims for osteoporosis medications within 1 year prior to fracture event.	Total, EasyCare and EasyCare Plus
Osteoporosis Screening in Older Women (OSW)	66 to 75 years of age	Female members between 65-75 years of age, who have been assessed for osteoporosis during the MY and the year prior.	ICD 10 Codes: NA CPT Codes: 76977, 77078, 77080, 77081, 77085 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) and advanced illness (A81.00) diagnosis, and members who had encounters for osteoporosis therapy during the MY or claims for osteoporosis medications within 3 years prior to the MY.	Total, EasyCare and EasyCare Plus

Effectiveness of Care- *Behavioral Health Measures:*

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Diagnoses of	1 years of	Members 1 years of age and older, who have	ICD 10 Codes: F03.90, F21, F31.0, F43.10	SelectHealth,
Mental Health	age and	been diagnosed with a mental health disorder	CPT Codes: NA	Total,
Disorders (DMH)	older	during the MY	Exclusions: Members on/using hospice services	EasyCare and
		Age stratifications are 1-17, 18-64 and 65 years of age and older.	(99377) and members who expired during the MY.	EasyCare Plus
Antidepressant	18 years of	Members 18 years of age and older with a	ICD 10 Codes: F32.0-F32.4, F32.9, F33.0-F33.3	SelectHealth,
Medication	age and	diagnosis of major depression, who received	CPT Codes: 99221, 98960, 90791, 98966	Total,
Management	older	and remained on antidepressant medications	Exclusions: Members on/using hospice services	EasyCare and
(AMM)		Rates reported include the effective acute	(99377), and members who expired during the	EasyCare Plus
		phase- those who received medication for at	MY.	
		least 84 days, and the effective continuation		
		phase- those who received medication for at		
		least 180 days.		
Follow-up Care for	6 to 12	Members between 6 to 12 years of age, who	ICD 10 Codes: F90.0-F90.2, F90.8, F90.9	SelectHealth
Children Prescribed	years of age	were prescribed ADHD (attention	CPT Codes: 96150-96154, 98960-98962	
ADHD Medication		deficit/hyperactivity disorder) medications	Exclusions: Members on/using hospice services	
(ADD)		and had a follow-up visit within 30 days of	(99377), members who expired during the MY,	
		prescription (initiation phase) and at least two	and members with a diagnosis of narcolepsy	
		more follow-up visits within 10 months of	(G47.411).	
		prescription (continuation and maintenance		
		phase).		
Follow-Up After	6 years of	Members 6 years of age and older, who had a	ICD 10 Codes: T14.91XA, F20.0, F31.0, F32.9	SelectHealth,
Hospitalization for	age and	mental illness or self-harm diagnosed hospital	CPT Codes: 98960-98962, 99201-99205	Total,
Mental Illness	older	admission, which was discharged and had a	Exclusions: Members on/using hospice services	EasyCare and
(FUH)		mental health provider follow-up. Follow-up	(99377), members who expired during the MY,	EasyCare Plus
		appointments are rated by those who	and any members discharged and readmitted	
		complete an appointment within 7 days post-	within 30 days.	
		discharge, and those who complete an		
		appointment within 30 days post-discharge.		

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Follow-Up After	6 years of	Members 6 years of age and older, who had a	ICD 10 Codes: T14.91XA, F20.0, F31.0, F32.9	SelectHealth,
Emergency	age and	mental illness or self-harm diagnosed	CPT Codes: 98960-98962, 99201-99205	Total,
Department Visit	older	emergency department (ED) visit, which was	Exclusions: Members on/using hospice services	EasyCare and
for Mental Illness		discharged and had a mental health provider	(99377), members who expired during the MY,	EasyCare Plus
(FUM)		follow-up. Follow-up appointments are rated	and members seen in the emergency	
		by those who complete an appointment	department resulting in a hospital admission or	
		within 7 days post ED-discharge, and those	an admission within 30 days of initial	
		who complete an appointment within 30 days	emergency department visit regardless of	
		post ED-discharge.	admission diagnosis.	
Diagnosed	13 years of	Members 13 years of age and older with a	ICD 10 Codes: F10.10, F11.10, F14.10, F15.10	SelectHealth,
Substance Use	age and	substance use disorder during the MY.	CPT Codes: NA	Total,
Disorders (DSU)	older	Substance abuse disorders include alcohol,	Exclusions: Members on/using hospice services	EasyCare and
		opioid, substance use and other/unspecified	(99377) and members who expired during the	EasyCare Plus
		drugs.	MY.	
Follow-Up After	13 years of	Members 13 years of age and older with a	ICD 10 Codes: F10.10, F11.10, F14.10, F15.10	SelectHealth,
High-Intensity Care	age and	substance use disorder, which had an acute	CPT Codes: 98960-98962	Total,
for Substance Use	older	inpatient hospitalization, residential	HCPCS Codes: H0017-H0019, H0008-H0014	EasyCare and
Disorder (FUI)		treatment, or detoxification visit, resulting in	Exclusions: Members on/using hospice services	EasyCare Plus
		follow-up visits. Follow-up appointments are	(99377), members who expired during the MY,	
		rated by those who complete an appointment	any nonacute inpatient admissions other than	
		within 7 days post-discharge, and those who	behavioral health.	
		complete an appointment within 30 days post-		
		discharge.		
Follow-Up After	13 years of	Members 13 years of age and older that had	ICD 10 Codes: T40.0X1A, F10.10, F11.10,	SelectHealth,
Emergency	age and	an emergency department visit with a primary	F14.10, F15.10	Total,
Department Visit	older	substance use disorder or drug overdose	CPT Codes: 98960-98962	EasyCare and
for Substance Use		diagnosis and completed a follow-up visit.	HCPCS Codes: H0020, H0004, G2075	EasyCare Plus
(FUA)		Follow-up appointments are rated by those	Exclusions: Members on/using hospice services	
		who complete an appointment within 7 days	(99377), members who expired during the MY,	
		post-discharge, and those who complete an	and members seen in the emergency	
		appointment within 30 days post-discharge.	department resulting in a hospital admission or	

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
			an admission within 30 days of initial emergency department visit regardless of admission diagnosis.	
Pharmacotherapy for Opioid Use Disorder (POD)	16 years of age and older	Members 16 years of age and older with an opioid use disorder, who received new pharmacotherapy that lasted for at least 180 days.	ICD 10 Codes: F11.10, F11.20, CPT Codes: NA HCPCS Codes: G2070, G2069, J0572, H0033, H0020, J2315 Exclusions: Members on/using hospice services (99377), members who expired during the MY, and members who had an inpatient stay for 8 or more days during treatment period.	SelectHealth, Total, EasyCare and EasyCare Plus
Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD)	18 to 64 years of age	Members between 18 and 64 years of age, diagnosed with schizophrenia, schizoaffective or bipolar disorder with a completed diabetes screening, which were prescribed antipsychotic medications during the MY.	ICD 10 Codes: F31.9, F20.9, F25.9 CPT Codes: 80047, 80053, 83036, 83037 Exclusions: Members who are diagnosed with diabetes or received insulin or oral hypoglycemic and/or antihyperglycemic medications during the MY or the prior year, and members that did not receive antipsychotic medications during the MY.	SelectHealth
Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD)	18 to 64 years of age	Members between 18 and 64 years of age, with a diagnosis of diabetes and schizophrenia or schizoaffective disorder, who completed an HbA1c and LDL-C test during the MY.	ICD 10 Codes: E10.9, E11.9, F20.9, F25.9 CPT Codes: 83036, 83037, 80061, 83700 CPT II Codes: 3044F, 3051F, 3052F, 3046F, 3048F, 3049F, 3050F Exclusions: Members on/using hospice services (99377), members who expired during the MY, members without a diagnosis of diabetes and diagnosed with polycystic ovarian syndrome (E28.2), gestational (O24.419) or steroid (E09.9) induced diabetes during the MY or the prior year.	SelectHealth

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC)	18 to 64 years of age	Members between 18 and 64 years of age, with a diagnosis of cardiovascular disease and schizophrenia or schizoaffective disorder, which completed LDL-C testing during the MY.	ICD 10 Codes: I20.0, I24.9, I21.4, F20.9, F25.9 CPT Codes: 80061, 83700, 83721 CPTII Codes: 3048F, 3049F, 3050F Exclusions: Members on/using hospice services (99377) and members who expired during the MY.	SelectHealth
Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA)	18 years of age and older	Members 18 years of age and older diagnosed with schizophrenia or schizoaffective disorder, which were prescribed an antipsychotic medication, and remained on treatment for at least 80% of the period.	ICD 10 Codes: F20.3, F20.89, F20.9, F25.8, F25.9 CPT Codes: NA HCPCS Codes: J2794, J0401, J1631, J1944, J2798, J2426m J2680, J2798 Exclusions: Members on/using hospice services (99377), members who expired during the MY, members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) and advanced illness (A81.00) diagnosis, members 81 and older with a frailty (L89.000) diagnosis, members with a diagnosis of Dementia (F0390) or do not have at least two antipsychotic prescription refill events.	SelectHealth, Total, EasyCare and EasyCare Plus
Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)	1 to 17 years of age	Members between 1 and 17 years of age, who were prescribed two or more antipsychotic medications and had metabolic testing. Metabolic testing includes blood glucose testing, cholesterol testing or both.	ICD 10 Codes: NA CPT Codes: 82465, 82947, 83036, 80061 CPTII Codes: 3044F, 3046F, 3051F, 3052F, 3048F, 3049F, 3050F Exclusions: Members on/using hospice services (99377) and members who expired during the MY.	SelectHealth
Utilization of the PHQ-9 to Monitor Depression Symptoms for	12 years of age and older	Members 12 years of age and older, that had an outpatient visit with a PHQ-9 assessment and score noted in the visit medical record,	ICD 10 Codes: NA CPT Codes: 99201-99205 Exclusions: Members on/using hospice services (99377), and members diagnosed with bipolar	SelectHealth, Total, EasyCare and EasyCare Plus

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Adolescents and Adults (DMS-E)		who are diagnosed with major depression or dysthymia.	disorder (F31.9), personality disorder, psychotic disorder or pervasive developmental disorder (F84.9).	
Depression Remission or Response for Adolescents and Adults (DRR-E)	12 years of age and older	Members 12 years of age and older, who are diagnosed with depression and have a high PHQ-9 score, that had successful remission (score <5) or response (score decreased by 50%) within 4 to 8 months after initial high score.	ICD 10 Codes: NA CPT Codes: 99201-99205 Exclusions: Members on/using hospice services (99377), and members diagnosed with bipolar disorder (F31.9), personality disorder, psychotic disorder or pervasive developmental disorder (F84.9).	SelectHealth, Total, EasyCare and EasyCare Plus

Effectiveness of Care- *Care Coordination Measures:*

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Advance Care Planning (ACP)	66 years of age and	Members between 66 and 80 years of age who have an advanced illness or frailty diagnosis, or	ICD 10 Codes: M62.81, Z74.01, G30.0, G30.9 CPT Codes: 99483, 99497	Total, EasyCare and
	older	receiving palliative care, and members 81 years of age and older who have received advanced care planning in the MY.	Exclusions: Members on/using hospice services (99377) and members who expired during the MY.	EasyCare Plus
Transitions of Care	18 years of	Members 18 years of age and older, who have	ICD 10 Codes: NA	Total,
(TRC)	age and	been discharged after an in-patient	CPT Codes: 99483, 99495, 98966, 99201	EasyCare and
	older	hospitalization, and have the following noted	Exclusions: Members on/using hospice services	EasyCare Plus
		in their outpatient medical record:	(99377) and members who expired during the	
		 Notification of Inpatient Admission 	MY.	
		within 2 days of discharge.		
		Receipt of Discharge information		
		within 2 days of discharge.		
		3. Patient Engagement within 30 days of		
		discharge. (cannot be completed the		
		same day of discharge)		
		4. Medication Reconciliation within 30		
		days of discharge. *Providers should		
- 11		use CPTII code 1111F.		
Follow-Up After	18 years of	Members 18 years of age and older, diagnosed	ICD 10 Codes: J44.9, G30.9, N18.9, I50.9, I21.9,	Total,
Emergency	age and	with multiple high-risk chronic conditions, who	148.91, 163.9	EasyCare and
Department Visit for	older	have been in the emergency department (ED)	CPT Codes: 99495, 98966, 99201, 90791	EasyCare Plus
People with		and completed a follow-up appointment	Exclusions: Members on/using hospice services	
Multiple High-Risk		within 7 days of the ED visit.	(99377) and members who expired during the	
Chronic Conditions		High-risk conditions include, but not limited to,	MY.	
(FMC)		COPD, Alzheimer's disease, chronic kidney		
		disease, heart failure, acute myocardial		
		infarction, atrial fibrillation, and stroke.		

HIV Care

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Viral Load Suppression	2 years of age and older	Members 2 years of age and older, who are diagnosed with HIV (Human Immunodeficiency Virus), with their last HIV Viral Load Testing of the year showing a viral load less than two hundred copies/mL.	ICD 10 Codes: NA CPT Codes: NA Exclusions: NA	SelectHealth
Sexually Transmitted Diseases: Chlamydia Screening Sexually Transmitted Diseases: Gonorrhea Screening	13 years of age and older 13 years of age and older	Members 13 years of age and older, who are diagnosed with HIV (Human Immunodeficiency Virus) who have completed a chlamydia screening within the MY. Members 13 years of age and older, who are diagnosed with HIV (Human Immunodeficiency Virus) who have completed a gonorrhea screening within the MY.	ICD 10 Codes: NA CPT Codes: NA Exclusions: Members under 18 years of age, who are not sexually active. ICD 10 Codes: NA CPT Codes: NA Exclusions: Members under 18 years of age, who are not sexually active.	SelectHealth SelectHealth
Sexually Transmitted Diseases: Syphilis Screening	18 years of age and older, as well as <18 who are sexually active	Members who are sexually active and/or over the age of 18, who are diagnosed with HIV (Human Immunodeficiency Virus) who have completed a serologic syphilis screening within the MY.	ICD 10 Codes: NA CPT Codes: NA Exclusions: Members under 18 years of age, who are not sexually active.	SelectHealth

Health Plan Descriptive Information

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Social Need	All ages	All members who have been screened for	ICD 10 Codes: NA	SelectHealth,
Screening and		Food, Housing and Transportation	CPT Codes: 96156, 96160, 96161	Total,
Intervention (SNS-E)		insecurity/needs, as well as have documented	Exclusions: Members on/using hospice services	EasyCare and
		interventions within 30 days of a positive	(99377) and members who expired during the	EasyCare Plus
		screening. Screenings should be completed at	MY, and members over 66 years of age residing	
		least annually.	in a facility.	
Language Diversity of	All ages	All members who have been screened for	ICD 10 Codes: NA	SelectHealth,
Membership (LDM)		their preferred spoken and written languages,	CPT Codes: NA	Total,
		which have been enrolled in the Health Plan	Exclusions: NA	EasyCare and
		at any time during the MY.		EasyCare Plus
Race/Ethnicity	All ages	Document and report the race and ethnicity	ICD 10 Codes: NA	SelectHealth,
Diversity of		of all members, enrolled in the Health Plan at	CPT Codes: NA	Total,
Membership (RDM)		any time during the MY.	Exclusions: NA	EasyCare and
				EasyCare Plus

Access/Availability of Care Measures:

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Adults' Access to	20 years	Members 20 years of age and older, who have	ICD 10 Codes: NA	SelectHealth,
Preventative/Ambulatory	of age	had a preventative care visit within the MY.	CPT Codes: 99201-99205, 99345	Total,
Health Services (AAP)	and	Preventative visits include ambulatory, online	Exclusions: Members on/using hospice	EasyCare
	older	and telephone visits.	services (99377) and members who expire	and
			during the MY.	EasyCare
				Plus
Initiation and	13 years	Members 13 years of age and older, who have a	ICD 10 Codes: F10.10, F11.10, F15.10	SelectHealth,
Engagement of	of age	new alcohol, opioid or other substance abuse	CPT Codes: 99408, 98960, 99212	Total,
Substance Use Disorder	and	disorder, which initiated and obtained treatment.	HCPCS Codes: G2086, G2075, G2080	EasyCare
Treatment (IET)	older	Initiation of SUD treatment completed within 14	Exclusions: Members on/using hospice	and
		days and engagement of SUD treatment	services (99377), members who expired	EasyCare
		completed within 34 days of initiation.	during the MY and any members where	Plus
			the initiation took place in an inpatient	
			admission where the member was	
			discharged after November 27 th of MY.	
Prenatal and Postpartum	None	Members who have had live births on October 8 th	ICD 10 Codes: O26.90, O26.91, O26.92,	SelectHealth
Care (PPC)	specified	the year prior through October 7 th of the MY.	O26.93	
		Measure includes timeliness of prenatal care	CPT Codes: 99201-99205, 99500, 59400,	
		during the first trimester, as well as postpartum	57170, 58300, 98966, 98969	
		care between 7 and 84 days post-delivery	Exclusions: Members on/using hospice	
		Members who have multiple live births during	services (99377), members who expired	
		one pregnancy count once, however multiple	during the MY, any non-live births and any	
		pregnancies resulting in two separate live	services provided in an acute inpatient	
		deliveries are counted twice.	setting (99221).	
Use of First-Line	1 to 17	Members between 1 and 17 years of age who	ICD 10 Codes: F20.9, F25.9, F33.3, F22	SelectHealth
Psychosocial Care for	years of	were prescribed a new antipsychotic medication,	CPT Codes: 98969-98972, 98966-98968,	
Children and Adolescents	age	with documented first-line psycho-social care.	Exclusions: Members on/using hospice	
on Antipsychotics (APP)			services (99377), members who expired	
			during the MY and members with	
			antipsychotic medication as first-line care.	

Measures Collected Through the Health Outcomes Survey:

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Medicare Health	65 years	This survey provides an idea of how well a	ICD 10 Codes: NA	Total,
Outcomes Survey (HOS)	of age	Medicare Advantage Organization manages	CPT Codes: NA	EasyCare and
	and	their members mental and physical health	Exclusions: NA	EasyCare Plus
	older	over a two year period. The rates of health-		
		status are indicated as better, same and		
		worse than expected.		
Fall Risk Management	65 years	Members who are 65 years of age and older,	ICD 10 Codes: NA	Total,
(FRM)	of age	who have seen a practitioner in the past 12	CPT Codes: NA	EasyCare and
	and	months and have discussed falls or	Exclusions: Members on/using hospice	EasyCare Plus
	older	gait/balance problems, and those who have	services (99377) per CMS administrative	
		fallen or had gait/balance problems in the	records.	
		last month that have received education on		
		prevention and treatment from their		
		practitioner.		
Management of Urinary	65 years	Members who are 65 years of age and older,	ICD 10 Codes: NA	Total,
Incontinence in Older	of age	that have reported urine leakage problems	CPT Codes: NA	EasyCare and
Adults (MUI)	and	with their health care providers in the past 6	Exclusions: Members on/using hospice	EasyCare Plus
	older	months, who have discussed the following	services (99377) per CMS administrative	
		with their health care provider: urinary	records.	
		incontinence, available treatment choices		
		and the impact urinary incontinence had on		
		daily living and sleep.		
Physical Activity in Older	65 years	Members who are 65 years of age and older,	ICD 10 Codes: NA	Total,
Adults (PAO)	of age	who have seen their doctor and discussed	CPT Codes: NA	EasyCare and
	and	their physical activity level, and those who	Exclusions: Members on/using hospice	EasyCare Plus
	older	received advice on starting, increasing or maintaining it, during the MY.	services (99377) per CMS administrative records.	

Utilization and Risk Adjusted Utilization Measures:

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Well-Child Visits in the First 30 Months of Life (W30)	15 to 30 months of age	Members between 31 days old to 15 months of age, who have had 6 or more well-child PCP visits within 15 months, as well as 2 or more well-child PCP visits between 15-30 months during the MY.	ICD 10 Codes: Z00.00, Z00.121, Z00.2, Z00.3 CPT Codes: 99381-99385, 99391-99395 Exclusions: Members on/using hospice services (99377) and members who expire during the MY.	SelectHealth
Child and Adolescent Well-Care Visits (WCV)	3 to 21 years of age	Members between 3 and 21 years of age, who have had one or more well-visits with their PCP or OB/GYN during the MY.	ICD 10 Codes: Z00.00, Z00.121, Z00.2, Z00.3 CPT Codes: 99381-99385, 99391-99395 Exclusions: Members on/using hospice services (99377) and members who expire during the MY.	SelectHealth
Plan All-Cause Readmissions (PCR)	18 years of age and older	Members who are 18 years of age and older, that have had an unplanned readmission, within 30 days of an observation event or inpatient admission during the MY.	ICD 10 Codes: NA CPT Codes: 99221-99223, 99217-99219 Exclusions: Members on/using hospice services (99377), members who expired during the hospitalization, and members who have a diagnosis of pregnancy or the discharge claim is for perinatal conditions.	SelectHealth, Total, EasyCare, and EasyCare Plus
Hospitalization Following Discharge from a Skilled Nursing Facility (HFS)	65 years of age and older	Members who are 65 years of age and older, who have transitioned from a Skilled Nursing Facility to the community and have had an unplanned inpatient admission within 30 to 60 days post transition.	ICD 10 Codes: NA CPT Codes: 99221-99223, 99217-99219 Exclusions: Members on/using hospice services (99377) and members residing in an institution long-term during the MY.	Total, EasyCare and EasyCare Plus
Acute Hospitalization Utilization (AHU)	18 years of age and older	The ratio for risk-adjustment of observed to expected inpatient admissions and observation discharges during the MY, for members who are 18 years of age and older.	ICD 10 Codes: NA CPT Codes: 99221-99223, 99217-99219 Exclusions: Members on/using hospice services (99377).	Total, EasyCare and EasyCare Plus

Emergency	18 years of	The ratio for risk-adjustment of observed to	ICD 10 Codes: NA	Total,
Department	age and	expected emergency department visits	CPT Codes: 99281-99285	EasyCare and
Utilization (EDU)	older	during the MY, for members who are 18	Exclusions: Members on/using hospice services	EasyCare
		years of age and older.	(99377), members with encounters for	Plus
			diagnosis of mental health or chemical	
			dependency, electroconvulsive therapy and	
			psychiatry.	
Hospitalization for	67 years of	Discharge rates for ambulatory care	ICD 10 Codes: NA	Total,
Potentially	age and	sensitive conditions per 1,000 members and	CPT Codes: 99221-99223, 99304-99310	EasyCare and
Preventable	older	the ratio for risk-adjustment of observed to	Exclusions: Members on/using hospice services	EasyCare
Complications (HPC)		expected discharges for acute and chronic	(99377), members residing in an institution	Plus
		ambulatory care sensitive conditions, for	long-term, and members enrolled in an	
		members who are 67 years of age and	institutional SNP during the MY.	
		older.		
Emergency	67 years of	The ratio for risk-adjustment of observed to	ICD 10 Codes: E10.64, E11.64, E13.64, E16.2	Total,
Department Visits for	age and	expected emergency department visits	CPT Codes: 99281-99285	EasyCare and
Hypoglycemia in	older	related to hypoglycemia during the MY.	Exclusions: Members on/using hospice services	EasyCare
Older Adults with		Rates are reported for members 67 years of	(99377), and members without a diagnosis of	Plus
Diabetes (EDH)		age and older with types 1 and 2 Diabetes,	diabetes and diagnosed with polycystic ovarian	
		with a subset of those with one or more	syndrome (E28.2), gestational (O24.419) or	
		insulin dispensing events within a 6-month	steroid (E09.9) induced diabetes.	
		treatment period July 1 the prior year		
		through December 31 of the MY.		

References and Comments:

- * HEDIS ® is a registered trademark of the National Committee for Quality Assurance (NCQA). Refer to www.ncqa.org/hedis/
- ❖ CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association
- * HCPCS: Healthcare Common Coding Procedure Coding System Level II
- LOINC (Logical Observation Identifiers Names and Codes)
- ❖ ICD-10: Centers for Medicare & Medicaid Services at www.cms.gov/Medicare/Coding/icd10/
- For additional information on Advanced Illness and Frailty, refer to https://www.ncqa.org/blog/improving-care-advanced-illness-

frailty/

Quality Assurance Reporting Requirements (QARR) are adopted from NCQA HEDIS® with New York state-specific measures added to address health issues of particular importance in New York. This document is not inclusive of all codes and measures. The measure descriptions and codes in this document are derived from the MY2023 HEDIS® Technical Specifications/Value Set Directory, and the Quality Assurance Reporting Requirements Technical Specifications manual. For the full list of codes, please refer to the NCQA store to purchase.



Have questions?

Please contact us at QualityManagement@vnshealth.org or Provider Education@vnshealth.org

For additional information and resources, please visit our provider website and toolkit at https://www.vnshealthplans.org/for-health-professionals-overview/